

AD-766 956

ON THE THERMAL LAG IN THE OCEAN DURING A  
PERIOD OF CLIMATIC CHANGE

I. I. Schell, et al

Ocean-Atmosphere Research Institute

Prepared for:

Air Force Office of Scientific Research

August 1973

DISTRIBUTED BY:

**NTIS**


National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE  
5285 Port Royal Road, Springfield Va. 22151

# On The Thermal Lag In The Ocean During A Period Of Climatic Change

By I.I. Schell and D.A. Corkum

Ocean-Atmosphere Research Institute

1640 Massachusetts Ave.  
Cambridge, Mass. 02138



D D C

AUG 31 1973

RECEIVED

DISSEMINATION STATEMENT A

Approved for public release  
Distribution Unlimited

Project OARI-1-73

Submitted as Annual Report to the Advanced Research Project Agency, ARPA Order No. 2159, Air Force  
Office of Scientific Research, under Contract Number F44620 72 C 0040

Technical Report: OARI-1-73

August 1973

NATIONAL TECHNICAL  
INFORMATION SERVICE  
11 Springfield Road  
Springfield, VA 22151

AD 766956

Unclassified  
Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Ocean-Atmosphere Research Institute 1640 Massachusetts Avenue Cambridge, Massachusetts 02138		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP ---	
3. REPORT TITLE  ON THE THERMAL LAG IN THE OCEAN DURING A PERIOD OF CLIMATIC CHANGE			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Annual Technical Report - June 1, 1972 - June 30, 1973			
5. AUTHOR(S) (First name, middle initial, last name)  I. I. Schell and D. A. Corkum			
6. REPORT DATE August 1973	7a. TOTAL NO. OF PAGES 2228	7b. NO. OF REFS 16	
8a. CONTRACT OR GRANT NO. F44620-72-C-0040	8b. ORIGINATOR'S REPORT NUMBER(S) ---		
8c. PROJECT NO. A02159			
8d. 62701D	9. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) --- AFOSR TR 73-1566		
10. DISTRIBUTION STATEMENT  Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES ---		12. SPONSORING MILITARY ACTIVITY AF Office of Scientific Research/NPG 1400 Wilson Boulevard Arlington, Va. 22209	
13. ABSTRACT An analysis was made of the sea surface temperatures in the northern No. Atlantic, sea temperature in the 0-200 m layer along the Kola meridian (70°E, approximately) between 70°30' and 72°30'N reflecting the inflow of relatively warm No. Atlantic water into the Barents Sea, and the sea surface temperatures in the Western Mediterranean affected by storms crossing from the No. Atlantic by the way of the Bay of Biscay, France and Spain and others which develop locally in wake of cold air outbreaks from the Continent; the ice off Iceland; and the air temperatures and pressures over the northern No. Atlantic, Greenland, and Europe during the period 1921-50 and the following decades 1951-60 and 1961-70, with the objective of determining a possible lag in the ocean vis-a-vis the atmosphere during a period of climatic change. The results obtained show a rise in the sea temperature in all three regions in the decade 1951-60 and with little change in the ice off Iceland from the period 1921-50 which preceded it, while the air temperatures were declining over the northern No. Atlantic indicating a lag in the ocean vis-a-vis the atmosphere. The results further show a fall in sea and air temperatures and sharp increase in the ice off Iceland in the decade 1961-70. One would expect should the decade 1971-80 show a rise in air temperatures, the sea temperature would lag less behind, since the warming would be limited to the surface layers primarily.			

DD FORM 1 NOV 65 1473

Security Classification

00  
11

Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Climatic Change Thermal Lag of Oceans Sea Temperatures Northern No. Atlantic Western Mediterranean Ice Off Iceland Air Temperatures						


Security Classification

ON THE THERMAL LAG IN THE OCEAN DURING A PERIOD  
OF CLIMATIC CHANGE

By I. I. Schell and D. A. Corkum

OCEAN-ATMOSPHERE RESEARCH INSTITUTE

1640 Massachusetts Ave.  
Cambridge, Mass. 02138



Project OARI-1-7,

Submitted as Annual Report to the Advanced Research Project  
Agency, ARPA Order No. 2159, Air Force Office of Scientific  
Research, under Contract Number F44620-72-C-0040

Technical Report: OARI-1-7,

August 1973

iv

Ocean-Atmosphere Research Institute  
1640 Massachusetts Avenue  
Cambridge, Massachusetts 02138


ANNUAL TECHNICAL REPORT

I. I. Schell  
and  
D. A. Corkum

August 1973

Sponsored by  
Advanced Research Projects Agency  
ARPA Order No. 2159

ARPA Order No. 2159  
Program Code - F10  
Name of Contractor - Ocean-Atmosphere Research Institute  
Effective Date of Contract - June 1, 1972  
Contract Expiration Date - June 30, 1973  
Amount of Contract Dollars - \$63,399  
Contract Number - F44620-72-C-0040  
Principal Investigator - I. I. Schell  
Phone Number - 617-491-8552  
Short Title of Work: Climatic Changes - Past 1,000 Years



The views and conclusions contained in this document are those of the author and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Projects Agency or the U.S. Government.



V

ABSTRACT

An analysis was made of the sea surface temperatures in the northern No. Atlantic, sea temperature in the 0-200 m layer along the Kola meridian ( $70^{\circ}\text{E}$ , approximately) between  $70^{\circ}30'$  and  $72^{\circ}30'\text{N}$  reflecting the inflow of relatively warm No. Atlantic water into the Barents Sea, and the sea surface temperatures in the Western Mediterranean affected by storms crossing from the No. Atlantic by the way of the Bay of Biscay, France and Spain and others which develop locally in wake of cold air outbreaks from the Continent; the ice off Iceland; and the air temperatures and pressures over the northern No. Atlantic, Greenland, and Europe during the period 1921-50 and the following decades 1951-60 and 1961-70, with the objective of determining a possible lag in the ocean vis-a-vis the atmosphere during a period of climatic change. The results obtained show a rise in the sea temperature in all three regions in the decade 1951-60 and with little change in the ice off Iceland from the period 1921-50 which preceded it, while the air temperatures were declining over the northern No. Atlantic indicating a lag in the ocean vis-a-vis the atmosphere. The results further show a fall in sea and air temperatures and sharp increase in the ice off Iceland in the decade 1961-70. One would expect should the decade 1971-80 show a rise in air temperatures, the sea temperature would lag less behind, since the warming would be limited to the surface layers primarily.

/

On the Thermal Lag in the Ocean during a Period  
of Climatic Change

By I. I. Schell and D. A. Corkum  
Ocean-Atmosphere Research Institute  
Cambridge, Mass. 02138

A. Introduction.

As can be seen from the extensive literature on past climates and their changes, a great deal of study and thought has been devoted to this problem.

Whether we look to predictive theories of the general circulation to provide a solution of this problem or seek its solution from a study of the changes in the past by discovering their orderly sequence, or trace a connection between them and solar variability, volcanicity, and other possible causes, there is the need for establishing as accurately as possible the climates of the past both on land and in the sea, and their spatial and physical coherence along with a need for defining the special characteristics of the climatic elements that distinguish one from another:

One such characteristic of a major climatic element we wish to investigate is the response of the ocean as compared with the atmosphere during a period of climatic change.

The difference in the thermal and dynamic characteristics between the ocean as represented by sea surface temperatures, ocean currents, and sea ice cover (in the higher latitudes) on the one hand, and the atmosphere, as represented by air temperature and



winds on the other, leads us to expect a slower and hence later response of the ocean during a period of climatic change than in the atmosphere. The following is an analysis of a possible time-lag of the response of the ocean vis a vis the atmosphere from a consideration of the sea surface temperatures in the northern No. Atlantic and Western Mediterranean, the sea temperature in the layer 0-200 m along the Kola ( $70^{\circ}\text{E}$ ) meridian, the sea surface temperatures from 8 Ocean Weather Stations in different locations in the No. Atlantic, the ice off Iceland, and finally, the pressure distribution (atmospheric circulation or winds) and air temperatures during the period 1921-1970 that was characterized by a rise, then a fall in temperature.

#### B. Analysis.

Mitchell (1961, 1963), following Willett (1950), has shown that beginning with the 1940's, the average "world" air temperature has taken a downward turn while remaining well above the long-term average. The change from the 1940-9 to the 1950-9 decades came, he found, to  $-0.19^{\circ}\text{C}$  for the year as a whole, and to  $-0.25^{\circ}\text{C}$  for the winter season.

Similarly, Rodewald in a series of papers (1963, 1967, 1972a, 1972b) treated in detail the sea surface temperatures in the northern No. Atlantic observed at the OWS ships which he compared with the air temperatures and the pressure distribution, his latest results showing a marked decrease in the air and sea temperatures in the No. Atlantic-Arctic in the most recent decade, 1961-70, as compared with that which preceded it. Earlier Smed has shown (1949) an increase in the sea surface temperature in the

northern No. Atlantic between  $50^{\circ}$  and  $66^{\circ}$ N, approximately, that was especially marked in the 1930's.

### 1. Pressure Distribution.

For evidence about the changes in circulation and winds during the 1921-70 period, we have compared the pressure over the northern No. Atlantic, Greenland, and Europe during the 1921-50 period with that in the 1951-60 decade and next, the pressures in this decade with those in the 1961-70 decade.

Figure 1 shows an area of excess pressure over Greenland and the vicinity in the 1951-60 decade over the 1921-50 period, suggesting an increased outflow of cold air southward from Greenland and lower air temperatures over East Greenland, Iceland, northwestern Europe and the North Atlantic in this decade (Fig. 2).

Similarly, the area of excess of pressure in the 1961-70 decade over the 1951-60 decade extending eastward from Greenland to Scandinavia and beyond (Fig. 3) signifies a greater outflow of cold air from the Arctic in this decade (1961-70) than in the preceding one as is shown by the lower air temperatures (Fig. 4). Also, the area of negative air temperature differences is larger.

### 2. Sea Temperatures.

#### a. Northern No. Atlantic Areas A-N.

For analysis of possible trends in the sea surface temperatures of the northern No. Atlantic, we have made use of the mean monthly tabulations of deviations of sea surface temperature north of  $50^{\circ}$ N in areas A to N, as designated by Smed (1949-1970), limiting ourselves to the April-September values for which more information is available than for the other months. The results obtained from the comparison of the temperatures in the 1951-60




Figure 1. Difference (1950/1-1959/60 minus 1920/1-1949/50) December-February Pressure (mb).




Figure 2. Differences (1951-1960 minus 1921-1950) mean annual air temperatures ( $^{\circ}\text{C}$ ). Circles denote locations of stations.






Figure 3. Difference (1960/1-1969/70 minus 1950/1-1959/60) December-February Pressure (mb).




Figure 4. Differences (1961-70 minus 1951-60) mean annual air temperatures ( $^{\circ}\text{C}$ ). Circles denote locations of stations including OWS ships.



decade with those of the 1921-50 period (Fig. 5) show higher values (positive differences) in all but one area (B) in the 1951-60 decade than in the earlier period. The air temperatures showed a small decline (Fig. 2). The results of the comparison in the sea surface temperatures in this region between the 1951-60 and 1961-70 decades show a sharp drop (Fig. 6).

b. Western Mediterranean.

Another comparison seeking to establish a thermal lag in the ocean vis-a-vis the atmosphere was made by means of the mean January-March and July-September sea surface temperatures of the Western Mediterranean. This region in the colder half of the year is subjected to No. Atlantic storms crossing into the area by the way of the Bay of Biscay, Spain, and France, and by storms which develop locally with cold air outbreaks from the continent sweeping over the relatively warm sea. Figure 7 shows the temperature in the 1951-60 decade was higher than in the 1921-50 period while the air temperatures were declining over France, Spain and elsewhere (see Fig. 2). The fall in the mean January-March sea surface temperature in the decade 1961-70 from the previous decade (Fig. 8) would be expected to have resulted from a southward shift in storm track over the No. Atlantic (Fig. 4) and more frequent outbreaks of colder air from the continent leading in each instance to a cooling of the waters and to lower sea surface temperatures.

A similar trend as for the January-March sea surface temperatures, namely, higher July-September sea surface temperatures in the 1951-60 decade than in the 1921-50 period was found in most areas also in this quarter (Fig. 9) while the air temperatures,




Figure 5. Differences (1951-1960 minus 1921-1950) April-September Sea Surface Temperature Areas A-N and Annual 0-200 m Sea Temperature Kola Section ( $^{\circ}\text{C}$ ).




Figure 6. Differences (1961-1970 minus 1951-1960) April-September Sea Surface Temperature Areas A-N and Annual 0-200 m Sea Temperature Kola Section ( $^{\circ}\text{C}$ ).




Figure 7. Difference (1951-60 minus 1921-50) January-March Sea Surface Temperature ( $^{\circ}\text{C}$ ) in the Western Mediterranean.




Figure 8. Difference (1961-70 minus 1951-60) January-March Sea Surface Temperature ( $^{\circ}\text{C}$ ) in the Western Mediterranean.






Figure 9. Differences (1951-1960 minus 1921-1950) July-September sea surface temperature ( $^{\circ}\text{C}$ ) Western Mediterranean.



as we have seen, showed a small decrease (Fig. 2).

One suspects that the increase in the July-September sea surface temperature in the 1951-60 decade, as well as the decrease in the 1961-70 decade from the decade which preceded it (Fig. 10) was in part due to the persistence of the temperatures anomaly in the cold season (January-March) since few if any storms cross into the western Mediterranean from the No. Atlantic in the warm half of the year.

c. Kola Meridian Section Temperature (0-200 m).

Another comparison to demonstrate a possible lag in the ocean vis a vis the air was made from a consideration of the Kola Meridian Section annual temperatures in the 0-200 m layer between latitudes  $70^{\circ}30'N$  and  $75^{\circ}23'N$  along the  $70^{\circ}E$  or Kola meridian. The small increase of  $0.16^{\circ}C$  in the temperatures from the 1921-50 period to the 1951-60 decade (Fig. 5) is in keeping with the increase in the sea surface temperature in the area to the south (A-N) as is the decrease of  $0.48^{\circ}C$  in the 1961-70 decade from the 1951-60 decade, the increase in temperature in the 1951-60 decade showing a thermal lag also for the inflow of warm water into the Barents Sea.

d. Ocean Weather Stations, OWS.

An independent comparison of the sea surface temperatures in the northern No. Atlantic is available from the frequent observations taken at the OWS ships which began in the late 1940's. The decrease for the most part in the temperature at these stations from the 1951-60 to the 1961-70 decades (Fig. 11) corroborates the findings based on the analysis of temperatures in the areas A to N (Fig. 6).




Figure 10. Differences (1961-1970 minus 1951-1960) July-September sea surface temperature ( $^{\circ}\text{C}$ ) Western Mediterranean.




Figure 11. Differences (1961-1970 minus 1951-1960) mean annual sea surface temperature North Atlantic Ocean Weather Stations ( $^{\circ}\text{C}$ ).

e. Ice off Iceland.

The severity of the ice off Iceland (Koch, 1945; Eythorsson and Sigtryggsson, 1971) for which data in some detail go back to the mid-18th century and in more general form to the time of settlement of Iceland, may be next considered for an analysis of a lag between the ocean and the atmosphere.

The sharp increase both in the duration and the spread of the ice along the coasts of Iceland in the 1961-70 decade over the previous decade (Fig. 12) is consistent with the greater build-up of cold air over northern Greenland and to the eastward, as well as an assumed greater frequency of northerly winds in that decade than in the earlier decades. The significant increase in the ice during this decade would be due to a marked southward transport of ice from the main pack in the north, and also to the formation of more young ice as a result of the lower air temperatures. The indicated smaller increase in northerly winds in the 1951-60 decade over the 1921-50 period than in the 1961-70 decade presumably was insufficient to produce a marked increase in the ice transport and formation of young ice.

f. No. Atlantic Weather Types (British Isles).

As further evidence of climatic changes from the 1921-50 to the 1951-60 and 1951-60 to the 1961-70 periods in the northern No. Atlantic, we investigated the trends in the frequencies of British Isles Weather Types. A decreased frequency of westerly type of circulation reflects a less zonal flow and an increased meridional type of circulation and hence lower temperatures. Thus, the average frequency of the westerly weather types (zonal flow) was 27.6% in the period 1929-49 as compared with 24.7% in




Figure 12. Ice off Iceland (duration in weeks multiplied by the number of areas with ice along the coasts). (After Koch, 1945; and Eythorsson and Sigtryggsson, 1971.)

the decade 1950-59 and only 21.9% in the following decade, 1960-69 (Lamb, 1972).

C. Data.

The data used in this study came from various sources. The mean monthly sea surface temperature deviations of the areas A-N (north of 50°N) were taken from the Annal. Biol., Cons. Internl. Explor. Mer, Vols. 2-26 (1947-1970).

The sea temperatures in the layer 0-200 m along the Kola meridian were adapted from tabulation by Sleptsov-Shevlevich (1967) with those for more recent years, from the Annal. Biol., Cons. Internl. Explor. Mer, Vols. 22-26. The ice off Iceland values were taken from computations by L. Koch (1945) as extended by J. Eythorsson and H. Sigtryggsson (1971). The OWS air and sea surface temperatures were taken from Mariners Weather Log and in part were made available by the Meteorol. Office, Bracknell. Other sources were World Weather Records and Monthly Climatic Data.

In seeking to establish the thermal lag between the oceans and the atmosphere, we have limited our study to data from the open ocean since observations at coastal stations when the prevailing winds are from land, would partially reflect also the air temperatures.

D. Remarks.

Since the increase and decrease, respectively, in the annual sea temperature in the 0-200 m layer along the Kola meridian reflects largely the advection of relatively warm water into the Barents Sea from the southwest, we may conclude



that the increase and decrease, respectively, in the sea surface temperature in areas A to N and at the OWS stations were also primarily the result of advection (see also Midtun, 1969). It will be noted that the decline in air temperatures in the 1950's did not reach the western mediterranean. Had this been the case, the increase in the mediterranean mean sea surface temperature might have been smaller.

The thermal lag in the ocean during the climatic change period 1921-1970 as exemplified primarily by the increase in the sea surface temperature during the 1951-60 decade while the air temperature was declining, presupposes climatic lags for other periods as well, and with longer time lags in the oceans during longer climatic or climate periods (see Emiliani, 1955).

#### E. Acknowledgements.

We are indebted to Mr. E. N. Sabbagh, Head, Synoptic Laboratory of the Institute for valuable help in the preparation of this work.

We are also indebted to the meteorological services of Denmark, France, Federal Republic of Germany, Iceland, Norway, Portugal, Spain, Sweden, and U. K. for making available the air temperatures from their respective regions and to officials of the NOAA for the tabulations of mean monthly values of sea surface temperatures of the Mediterranean used in our study.

The work was carried out with support of the Advanced Projects Research Agency, Air Force Office of Scientific Research, USAF.

### References

- Dickson, R. R. and H. H. Lamb, 1971: A review of recent hydro-meteorological events in the No. Atlantic sector. Intern. Comm. Northwest Atlantic Fisheries Sp. Publ. No. 8. pp. 55-62.
- Eythorsson, J. and H. Sigtrygsson, 1971: The climates and weather of Iceland. In "The Zoology of Iceland." 1(3): 1-62. E. Munksgaard, (Copenhagen and Reykjavik).
- Emiliani, C., 1955: Pleistocene temperatures. J. Geol. 63: 538-70.
- Koch, L., 1945: The East Greenland Ice. Medd. Gronland 130(3): 373 pp.
- Lamb, H. H., 1972: British Isles Weather Types and a register of the daily sequence of circulation type patterns 1861-1971. Geophys. Mem. No. 116. Met. Office, London, 85 pp.
- Midtun, L., 1969: Variability of temperature and salinity at some locations off the coast of Norway. Prog. in Ocean 5: 41-54. Pergamon Press.
- Mitchell, J. M., Jr., 1961: Recent secular changes of global temperature. Ann. N. Y. Acad. Sci. 95(1): 255-50.
- \_\_\_\_\_, 1963: On the world-wide pattern of secular temperature change. In "Changes of Climate." Arid Zone Research, XX Proc. Rome Symp. UNESCO-WMO, pp 161-80.
- Rodewald, M., 1963: Sea-surface temperatures of the North Atlantic Ocean during the decade 1951-60, their anomalies and development in relation to the atmospheric circulation. In "Changes of Climate." Arid Zone Research, XX Proc. Rome Symp. UNESCO-WMO, pp. 107-97.
- \_\_\_\_\_, 1967: Recent variations of North Atlantic sea surface temperatures and the "type tendencies" of the atmospheric circulation. Intern. Comm. Northw. Atlant. Fish. Red-book, 1967, Part IV, pp. 6-23. Dartmouth, N. S.
- \_\_\_\_\_, 1972a: Temperature condition in the North and Northwest Atlantic during the decade 1961-70. Intern. Comm. Northw. Atlant. Fish. Sp. Publ. No. 8, pp. 9-34. Dartmouth, N. S.
- \_\_\_\_\_, 1972b: Long-term variations of sea temperature in the areas of nine north Atlantic Ocean Weather Stations during the period 1951-1968. Cons. Perm. Intern. Explor. Mer, Rapp. Proces Verb. 162: 139-153.
- Sleptsov-Shevlevich, B. A., 1967: Study of instability in helio-geophysical relationships. In "Ice Cover of the Arctic Ocean

and a Method for Calculating Certain Ice Phenomena." Trudy Arctic-Antarctic Sci. Res. Inst. No. 257. Also Naval Intel. Comm. Transl. No. 2641: 133-68. Washington, D. C.

Smed, J., 1949: The increase in the sea temperature in northern waters during recent years. Cons. Perm. Internl. Explor. Mer, Rapp. Proc. Verb. 125: 21-25.

\_\_\_\_\_, 1947-1970: Monthly anomalies of the surface temperature in sea. Ann. Biol. 2-26. Copenhagen.

Willetts, H. C., 1950: Temperature trends of the past century. In Centenary Proc. Royal Meteorol. Soc. pp. 195-206.